

Linear ODEs & Divergent Series

Marc Mezzarobba

Algorithmic Combinatorics seminar, RISC, 2013-06-05

Disclaimer

I am not an expert
...just trying to learn about this



Werner Balser

Formal Power Series and Linear Systems of Meromorphic Ordinary Differential Equations



Marius van der Put & Michael F. Singer

Galois Theory of Linear Differential Equations

Chap. 7: "Exact Asymptotics"

Singularities of Linear ODEs

$$a_r(z) y^{(r)}(z) + \cdots + a_1(z) y'(z) + a_0(z) y(z) = 0, \quad a_k \in \mathbb{C}[z]$$

$a_r(0) \neq 0$ — Ordinary point

$$y(z) = \sum_{n=0}^{\infty} y_n z^n \quad \text{analytic on } D(0, \rho)$$

$a_r(0) = 0$ — Singular point

$$y_i(z) = \sum_j (\log z)^j z^{\lambda_{i,j}} y_{i,j}(z) \quad y_{i,j} \text{ analytic on } D(0, \rho) \setminus \{0\}$$

- All $y_{i,j}(z)$ have a **pole** at 0:
regular singular point
- Some $y_{i,j}(z)$ has an **essential singularity**:
irregular singular point

Formal Solutions


$$y_i(z) = \underbrace{e^{\text{Poly}_i\left(\frac{1}{z^{1/q}}\right)} z^{\lambda_i}}_{\substack{\text{analytic in} \\ \text{suitable} \\ \text{domains}}} \sum_j^{\text{(finite)}} (\log z)^j \underbrace{\sum_{n=0}^{\infty} c_{i,j,n} z^{n/q}}_{\substack{\in \mathbb{C}[[z^{1/q}]] \\ \text{in general divergent}}},$$

Example

$$\hat{y}(z) = \sum_{n=0}^{\infty} n! z^n \quad \text{satisfies} \quad z^2 \hat{y}''(z) + (3z - 1) \hat{y}'(z) + \hat{y}(z) = 0$$

“True” solutions on $\mathbb{C} \setminus e^{i\theta} \mathbb{R}_+$:

$$\left\langle \frac{1}{z} e^{-1/z}, \quad \frac{1}{z} e^{-1/z} \operatorname{Ei}\left(\frac{1}{z}\right) \right\rangle$$




Approximation by Divergent Series

$$w(z) = z e^{-z} \operatorname{Ei}(z)$$

$$w(10) \approx 1.13147$$

$$\hat{w}_{;N}(z) = \sum_{n=0}^{N-1} \frac{n!}{z^n}$$


$$\hat{w}_{;10}(10) \approx 1.13159$$



Summation to the Least Term

N	$\hat{w}_{;N}(10)$
1	1.00000
2	1.10000
3	1.12000
4	1.12600
5	1.12840
6	1.12960
7	1.13032
8	1.13082
9	1.13122
10	1.13158
11	1.13194
12	1.13234
13	1.13282
14	1.13344
15	1.13431


$$w(10) \approx 1.13147$$



$\text{RelativeError}(N, 10)$


Summation to the Least Term

$$\hat{w}(10) = 1 + \frac{1}{10} + \frac{2}{10^2} + \cdots + \frac{8!}{10^8} + \frac{\mathbf{9!}}{\mathbf{10^9}} + \frac{\mathbf{10!}}{\mathbf{10^{10}}} + \frac{11!}{10^{11}} + \frac{12!}{10^{12}} + \cdots$$




Sectors and Regions

Sector




$$\left\{ z \in \dot{\mathbb{C}} : 0 < |z| < \rho_0, \theta_0 < \arg z < \theta_1 \right\}$$

Sectorial region of opening α



- contained in a sector of opening α
- contains sectors of opening $\alpha - \varepsilon$



$\dot{\mathbb{C}}$

$$0 < \rho < \infty$$
$$\theta \in \mathbb{R}$$

(Poincaré) Asymptotic Expansions

Given y analytic in G , we write

$$y(z) \cong \hat{y}(z) = \sum_{n=0}^{\infty} y_n z^n \in \mathbb{C}[[z]] \quad \text{as } z \rightarrow 0 \text{ with } z \in G$$

if **for all N** and for any closed sector $\Delta \subset G$,

$$y(z) = y_0 + y_1 z + \cdots + y_{N-1} z^{N-1} + O(z^N), \quad z \rightarrow 0, z \in \Delta$$

where the constant hidden in the $O(\cdot)$ may depend on N and Δ .

- A given y has at most one asymptotic expansion (its Taylor expansion!)
- Any \hat{y} is the asymptotic expansion of infinitely many functions

$$e^{-1/z} = 0 + 0 z + \cdots + 0 z^{N-1} + O(z^N) \quad \text{as } z \rightarrow 0, \operatorname{Re} z > \varepsilon$$

The Main Asymptotic Existence Theorem

Theorem [Poincaré, Horn, Birkhoff...]

In any sufficiently small sector Δ with apex at 0, there exists a basis of **analytic solutions** $y_i: \Delta \rightarrow \mathbb{C}$ with **asymptotic expansions**

$$y_i(z) \cong e^{\text{Poly}_i\left(\frac{1}{z^{1/q}}\right)} z^{\lambda_i} \sum_j^{\text{(finite)}} (\log z)^j \sum_{n=0}^{\infty} c_{i,j,n} z^{n/q}, \quad z \rightarrow 0, \quad z \in \Delta.$$



Wolfgang Wasow

Asymptotic Expansions for Ordinary Difference Equations

- Uniqueness?
- Computation?
- Sectors of validity?
- Link between different asymptotic lifts?


The Stokes Phenomenon

$$y''(z) + 2z y'(z) = 0$$

$$-\zeta^3 w''(\zeta) + 2(\zeta^2 - 1) w'(\zeta) = 0$$

Analytic solutions:


$$\langle 1, \operatorname{erfc} z \rangle$$




Formal solutions at ∞ :

$$\left\langle 1, e^{-z^2} \left(\frac{1}{z} - \frac{1}{2z^3} + \frac{3}{4z^5} - \dots \right) \right\rangle$$

$$\operatorname{erfc}(z) \cong \frac{e^{-z^2}}{\sqrt{\pi}} \left(\frac{1}{z} - \frac{1}{2z^3} + \frac{3}{4z^5} - \dots \right), \quad z \rightarrow \infty \quad \text{in}$$



$$\operatorname{erfc}(z) \cong 2 + \frac{e^{-z^2}}{\sqrt{\pi}} \left(\frac{1}{z} - \frac{1}{2z^3} + \frac{3}{4z^5} - \dots \right), \quad z \rightarrow \infty \quad \text{in}$$



The Stokes Phenomenon

$$\begin{aligned}z - \frac{1}{2} z^3 + \frac{3}{4} z^5 - \frac{15}{8} z^7 + \frac{105}{16} z^9 - \frac{945}{32} z^{10} + \dots + \\ \approx \sqrt{\pi} \exp\left(\frac{1}{z^2}\right) \left(\textcolor{blue}{A} \operatorname{erfc} \frac{1}{z} + \textcolor{blue}{B} 1 \right)\end{aligned}$$

Gevrey Series

Formal power series of Gevrey order $s > 0$:

$$\mathbb{C}[[z]]_s = \left\{ \sum_{n \geq 0} y_n z^n : y_n = O(\mathbf{n!}^s e^{O(n)}) \right\}$$

(= Taylor expansions of \mathcal{C}^∞ functions whose derivatives
“do not grow too fast”)

$$\begin{aligned} \mathbb{C}\{z\} &= \mathbb{C}[[z]]_0 \subseteq \mathbb{C}[[z]]_s \subsetneq \mathbb{C}[[z]]_{s'} \subsetneq \mathbb{C}[[z]] \end{aligned}$$

$s < s'$

Gevrey Asymptotics

Given y analytic on G , we write

$$y(z) \underset{s}{\cong} \hat{y}(z) = \sum_{n=0}^{\infty} y_n z^n \in \mathbb{C}[[z]] \quad \text{as } z \rightarrow 0 \text{ with } z \in G$$

if for all N and for any closed sector $\Delta \subset G$,

$$|y(z) - (y_0 + y_1 z + \cdots + y_{N-1} z^{N-1})| \leq c K^N N!^s$$


where c, K do not depend on N (but may depend on Δ).

$$y = \hat{y} \in \mathbb{C}\{z\} \Leftrightarrow y \underset{0}{\cong} \hat{y} \Rightarrow y \underset{s}{\cong} \hat{y} \Rightarrow y \underset{s'}{\cong} \hat{y} \Rightarrow y \cong \hat{y}$$


$$s < s'$$

Uniqueness

Narrow Regions



Wide Regions



Proposition ("Ritt's Theorem")

Let $G \subset \dot{\mathbb{C}}$ of opening $\leq s\pi$, and let $\hat{f} \in \mathbb{C}[[z]]_s$.

Then there exists an analytic function $f: G \rightarrow \mathbb{C}$ s.t.

$$f \cong_s \hat{f} \quad \text{in } G.$$

Such a function is never unique.

Proposition (Watson's Lemma)

Let $G \subset \dot{\mathbb{C}}$ of opening $> s\pi$, and let $f: G \rightarrow \mathbb{C}$ be an analytic fun s.t.

$$f \cong \hat{0} \quad \text{in } G.$$


Then $f \equiv 0$.

The mapping $y \mapsto$ its asymptotic expansion is never surjective.

The Laplace Transform

$$\mathcal{L}_k f(z) = z^{-k} \int_0^\infty f(u) e^{-(u/z)^k} d(u^k)$$

- f analytic around $(0, \infty)$, continuous at 0
- $f(u) = O(e^{cu^k})$ as $u \rightarrow \infty$, $u \in \mathbb{R}$
- $\mathcal{L}_k f$ is defined (at least) for $\operatorname{Re}(z^{-k}) > c$
i.e., in a region of opening π/k




$$\operatorname{Re}(z^{-2}) > 1$$

The Laplace Transform in a Sector

$$\mathcal{L}_k f(z) = z^{-k} \int_0^{e^{i\theta}\infty} f(u) e^{-(u/z)^k} d(u^k)$$

- Replace $(0, \infty)$ by $(0, e^{i\theta}\infty)$
⇒ Laplace transform on a sector



The Laplace Transform of a Monomial

$$\mathcal{L}_k f(z) = z^{-k} \int_0^\infty f(u) e^{-(u/z)^k} d(u^k)$$

Laplace transform of a monomial (assume $z > 0$)

$$\begin{aligned}\mathcal{L}_k (u^n)(z) &= z^{-k} \int_0^\infty u^n e^{-(u/z)^k} d(u^k) \\ &= \int_0^\infty z^n v^{n/k} e^{-v} dv & v = \left(\frac{u}{z}\right)^k \\ &= \Gamma\left(1 + \frac{n}{k}\right) z^n & d(u^k) = z^k dv\end{aligned}$$


The Formal Laplace Transform

$$\hat{\mathcal{L}}_k \hat{f}(z) := \sum_n \Gamma\left(1 + \frac{n}{k}\right) f_n z^n, \quad \hat{f} = \sum_n f_n u^n \in \mathbb{C}[[u]]$$

Property

In the setting of the previous slides


$$f \cong_s \hat{f} \implies \mathcal{L}_k f \cong_{s + \frac{1}{k}} \hat{\mathcal{L}}_k \hat{f}$$



The Borel Transform

$$\mathcal{B}_k f(u) = \frac{1}{2\pi i} \int_{\gamma} z^k f(z) e^{(\frac{u}{z})^k} d(z^{-k})$$

$$\hat{\mathcal{B}}_k \hat{f}(u) = \sum_{n=0}^{\infty} \frac{f_n}{\Gamma(1 + \frac{n}{k})} u^n$$



- f analytic on a region G of opening $\alpha > \pi/k$, bounded as $z \rightarrow 0$
- $\mathcal{B}_k f$ well-defined when $\operatorname{Re}((u/z)^k) < 0$ on the radial rays, i.e., $|\arg u| < \varepsilon$
- Rotate the integration path to define $\mathcal{B}_k f$ on a sector of opening $\alpha - \frac{\pi}{k}$
- $\mathcal{B}_k f(u) \lesssim e^{|u|^k}$ as $u \rightarrow \infty$ in that region
- Respects Gevrey asymptotics

Borel-Laplace for Convergent Series

- The (analytic) Borel and Laplace transforms of order k are inverse of each other
- In particular, for **convergent** series :

$$f(z) = \sum_n f_n z^n \in \mathbb{C}\{z\} \quad f(z) = \int_0^{\infty(d)} g(u) e^{-(u/z)^k} d(u^k)$$


$$\downarrow \mathcal{B}_k \qquad \qquad \uparrow \mathcal{L}_k$$

$$g(u) = z^{-k} \int_0^{\infty(d)} f(u) e^{-(u/z)^k} d(u^k) = \sum_n \frac{f_n}{\Gamma(1 + \frac{n}{k})} u^n$$

- entire function
- $\lesssim e^{-|u|^k}$


k -Summability

$$\hat{f}(z) = \sum_n f_n z^n \in \mathbb{C}[[z]]_{1/k} \quad \cong_{1/k} \quad f(z) = \int_0^{\infty(d)} g(u) e^{-(u/z)^k} d(u^k)$$



$$\downarrow \hat{\mathcal{B}}_k$$

$$\hat{g}(u) = \sum_n \frac{f_n}{\Gamma(1 + \frac{n}{k})} u^n \in \mathbb{C}\{u\}$$




analytic
continuation



“succeeds” iff
 $\exists f, \hat{f} \cong_{1/k} f$
 in a wide sector
 bisected by d

$$\uparrow \mathcal{L}_k$$

$$g(u)$$



Example

$$\hat{f}(z) = \sum_n n! z^n \in \mathbb{C}[[z]]_1$$

$$f(z) = z^{-1} \int_0^{\infty(d)} \frac{e^{-u/z}}{1-u} du$$

$\downarrow \hat{\mathcal{B}}_k$

$\uparrow \mathcal{L}_k$

$$\begin{array}{ccc} \hat{g}(u) = \sum_n u^n \in \mathbb{C}\{u\} & \xrightarrow{\text{an. cont.}} & g(u) = \frac{1}{1-u} \ll e^{|u|} \\ u \notin \mathbb{R}_+ & & \end{array}$$

E.g., with $d = e^{i \cdot \pi}$ and $z < 0$:

$$\begin{aligned} z^{-1} \int_0^{-\infty} \frac{e^{-u/z}}{1-u} du &= -z^{-1} \int_0^{-\infty} \frac{e^v}{1+zv} z dv = z^{-1} e^{-1/z} \int_{-\infty}^0 \frac{\exp(v+z^{-1})}{v+z^{-1}} dv \\ &= z^{-1} e^{-1/z} \int_{-\infty}^{-1/z} \frac{e^{-w}}{w} dw = z^{-1} e^{-1/z} \operatorname{Ei}(-1/z). \end{aligned}$$

Properties of the k -Summation Operator

$\mathbb{C}\{z\}_{k,d} \subsetneq \mathbb{C}[[z]]_{1/k}$: space of k -summable functions in the direction d
 $\mathcal{S}_{k,d}$: k -summation operator

$$\begin{aligned}\mathcal{S}_{k,d}(\alpha \hat{f}) &= \alpha \mathcal{S}_{k,d}(\hat{f}) & \mathcal{S}_{k,d}(\hat{f} \hat{g}) &= \mathcal{S}_{k,d}(\hat{f}) \mathcal{S}_{k,d}(\hat{g}) \\ \mathcal{S}_{k,d}(\hat{f} + \hat{g}) &= \mathcal{S}_{k,d}(\hat{f}) + \mathcal{S}_{k,d}(\hat{g}) & \mathcal{S}_{k,d}(\hat{f}') &= \mathcal{S}_{k,d}(\hat{f})'\end{aligned}$$

Hence:

$$\underbrace{L\left(z, \frac{d}{dz}\right) \cdot \hat{y}}_{\substack{\rightarrow \in \mathbb{C}\{z\}_{k,d} \\ \text{Linear differential operator}}} = 0 \Rightarrow L\left(z, \frac{d}{dz}\right) \cdot (\mathcal{S}_{k,d} \hat{y}) = 0$$

Linear differential operator

$$\in \mathbb{C}(x)\left\langle \frac{d}{dz} \right\rangle$$